

Immunization Schedule (IAP Recommended-2013)

AGE	DISEASE	VACCINE	DOSE	ROUTE	DUE ON	GIVEN ON
At Birth	Tuberculosis	BCG	-	Intradermal Injection		
	Polio	OPV (oral polio vaccine)	0	Oral Drops		
	Hepatitis B	Hep B	1 st	Intramuscular Injection		
6 Weeks (1.5 months)	Diphtheria + Tetanus + Pertusis	DTwP/DTaP	1 st	Intramuscular Injection		
	Haemophilus influenza type b	Hib	1 st	Intramuscular Injection		
	Rotavirus	Rotavirus	1 st	Oral Drops		
	Pneumococcal infections	PCV (Pneumococcal conjugate vaccine)	1 st	Intramuscular Injection		
	Polio	IPV (Injectable polio vaccine)	1 st	Intramuscular Injection		
	Hepatitis B	Hep B	1 st	Intramuscular Injection		
10 Weeks (2.5 months)	Diphtheria + Tetanus + Pertusis	DTwP/DTaP	2 nd	Intramuscular Injection		
	Haemophilus influenza type b	Hib	2 nd	Intramuscular Injection		
	Rotavirus	Rotavirus	2 nd	Oral Drops		
	Pneumococcal infections	PCV (Pneumococcal conjugate vaccine)	2 nd	Intramuscular Injection		
	Polio	IPV (Injectable polio vaccine)	2 nd	Intramuscular Injection		
14 Weeks (3.5 months)	Diphtheria + Tetanus + Pertusis	DTwP/DTaP	3 rd	Intramuscular Injection		
	Haemophilus influenza type b	Hib	3 rd	Intramuscular Injection		
	Rotavirus	Rotavirus	3 rd	Oral Drops		
	Pneumococcal infections	PCV (Pneumococcal conjugate vaccine)	3 rd	Intramuscular Injection		
	Polio	IPV (Injectable polio vaccine)	3 rd	Intramuscular Injection		
6 months	Polio	OPV (oral polio vaccine)	1 st	Oral Drops		
	Hepatitis B	Hep B	3 rd	Intramuscular Injection		
9 months	Polio	OPV (oral polio vaccine)	2 nd	Oral Drops		
	Mumps + Measles + Rubella	MMR	1	Subcutaneous Injection		
9 -12 Months	Typhoid	Typhoid Conjugate vaccine	1	Intramuscular Injection		
1 year	Hepatitis A	Hep A	1 st	Intramuscular Injection		
15 months (1 year and 3 months)	Mumps + Measles + Rubella	MMR	2 nd	Subcutaneous Injection		
	Chicken Pox	Varicella	1 st	Subcutaneous Injection		

months)	Pneumococcal infections	PCV (Pneumococcal conjugate vaccine)	Booster	Intramuscular Injection		
16 to 18 months	Diphtheria + Tetanus + Pertusis	DTwP/DTaP	Booster 1	Intramuscular Injection		
	Haemophilus influenza type b	Hib	Booster 1	Intramuscular Injection		
	Polio	IPV (Injectable polio vaccine)	Booster 1	Intramuscular Injection		
18 months	Hepatitis A	Hep A	2 nd	Intramuscular Injection		
2 years	Typhoid	Typhoid conjugate vaccine	Booster	Intramuscular Injection		
4 to 6 years	Diphtheria + Tetanus + Pertusis	DTwP/DTaP	Booster 2	Intramuscular Injection		
	Polio	OPV (oral polio vaccine)	3 rd	Oral Drops		
	Chicken Pox	Varicella	2 nd	Subcutaneous Injection		
	Typhoid	Typhoid	Booster	Intramuscular Injection		
10 to 12 years	Tetanus	Tdap/Td		Intramuscular Injection		
	Cervical Cancer (FEMALES ONLY)	HPV (Human Papilloma Virus)	3 doses in total	Intramuscular Injection		

FLU VACCINE:

Additionally to the above mentioned IAP schedule, paediatricians also recommend an influenza or **FLU** vaccine which protects from Influenza Vaccine (A&B), H1N1 Vaccine (Swine Flu).

Dosage recommendations for Flu vaccine:

Age	Dose	No. of doses (first vaccination)	No. of doses [#] (subsequent years)
6 months to 35 months	0.25 mL	2 [§]	1
1 3 to 9 years	0.5 mL	2 [§]	1
≥ 10 years	0.5 mL	1	1

[#]Where children 6 months to ≤ 9 years of age receiving influenza vaccine for the first time have not received the second dose within the same year, they should be given 2 doses the following year.

[§] For children aged ≤ 9 years who are receiving influenza vaccine for the first time, it is recommended that they receive 2 doses at least 1 month apart.

*** Please note, every paediatrician may follow different guidelines, depending upon the type of vaccines used. Hence detailed discussion with your doctor is advised.**

© Gauri Kshirsagar 2015